arXiv:1704.07151v1 [hep-ph] 24 Apr 2017

Can we measure θ_{23} octant in 3+1 scheme?

Sanjib Kumar Agarwalla[1,2,*], Sabya Sachi Chatterjee[1,2,†,§], and Antonio Palazzo[3,4,‡]

[1] Institute of Physics, Sachivalaya Marg, Sainik School Post, Bhubaneswar 751005, India

[2] Homi Bhabha National Institute, Training School Complex, Anushakti Nagar, Mumbai 400085, India

[3] Dipartimento Interateneo di Fisica "Michelangelo Merlin," Via Amendola 173, 70126 Bari, Italy

[4] Istituto Nazionale di Fisica Nucleare, Sezione di Bari, Via Orabona 4, 70126 Bari, Italy

[*] sanjib@iopb.res.in, [†] sabya@iopb.res.in, [‡] palazzo@ba.infn.it

Abstract. Current 3ν global fits predict two degenerate solutions for θ_{23}: one lies in lower octant ($\theta_{23} < \pi/4$), and the other belongs to higher octant ($\theta_{23} > \pi/4$). Here, we study how the measurement of θ_{23} octant would be affected in the upcoming Deep Underground Neutrino Experiment (DUNE) if there exist a light eV-scale sterile neutrino. We show that in 3+1 scheme, a new interference term in $\nu_\mu \to \nu_e$ oscillation probability can spoil the chances of measuring θ_{23} octant completely.

Keywords: Octant of θ_{23}, sterile neutrino, Long-Baseline experiments

Introduction: The resolution of octant[¶] [1] of θ_{23} is one of the fundamental problems in neutrino oscillation. Long-baseline (LBL) experiments [2] can resolve this octant ambiguity of θ_{23} with the help of $\nu_\mu \to \nu_e$ appearance channel, and the vital information coming from $\nu_\mu \to \nu_\mu$ disappearance channel also play an important role. Interestingly, at present, there are short-baseline anomalies which hint towards the existence of light eV-scale sterile neutrino [3]. Here, we expound in detail the capability of proposed LBL experiment DUNE to measure θ_{23} octant considering one light eV-scale sterile neutrino along with three active neutrinos.

Theoretical framework: In the 3+1 scheme, a new a mass eigenstate ν_4 appears on top of 3ν framework whose mixing is parametrized as

$$U = \tilde{R}_{34} R_{24} \tilde{R}_{14} R_{23} \tilde{R}_{13} R_{12}\,, \tag{1}$$

where R_{ij} ($\tilde{R}_{ij}$) is a real (complex) rotation in the (i,j) plane. The details of the parametrization of U can be seen in [4].

[§] Speaker, corresponding author.

[¶] According to the present 3ν best-fit [5], θ_{23} can have two solutions: one $< \pi/4$, labelled as lower octant (LO), and other $> \pi/4$, known as higher octant (HO).

In [4], it was shown that the 4-flavor appearance probability can be approximately expressed as the sum of three terms $P^{4\nu}_{\mu e} \simeq P_0 + P_1 + P_2$, which in vacuum appears as

$$P_0 \simeq 4 s^2_{23} s^2_{13} \sin^2 \Delta \,, \tag{2}$$

$$P_1 \simeq 8 s_{13} s_{12} c_{12} s_{23} c_{23} (\alpha \Delta) \sin \Delta \cos(\Delta \pm \delta_{13}) \,, \tag{3}$$

$$P_2 \simeq 4 s_{14} s_{24} s_{13} s_{23} \sin \Delta \sin(\Delta \pm \delta_{13} \mp \delta_{14}) \,. \tag{4}$$

where $\Delta \equiv \Delta m^2_{31} L/4E$ and $\alpha \equiv \Delta m^2_{21}/\Delta m^2_{31}$. In the double sign, the upper (lower) sign corresponds to neutrinos (antineutrinos). The new interference term P_2 is governed by the interference between the atmospheric frequency and the large frequency related to the new mass eigenstate [4] which gets averaged out by the finite energy resolution of the detector. Recent global fits [5,6,7] suggests $s_{13} \sim s_{14} \sim s_{24} \sim 0.15\,(\sim \epsilon)$ and $\alpha = 0.03\,(\sim \epsilon^2)$ implying $P_0 \sim \epsilon^2$, $P_1 \sim \epsilon^3$, $P_2 \sim \epsilon^3$. An experiment can measure the octant of θ_{23} even in the presence of unknown CP-phases, if there is a difference between the probabilities corresponding to the different octants, i.e.

$$\Delta P \equiv P^{\rm HO}_{\mu e}(\delta^{\rm HO}_{13}, \delta^{\rm HO}_{14}) - P^{\rm LO}_{\mu e}(\delta^{\rm LO}_{13}, \delta^{\rm LO}_{14}) \neq 0 \,, \tag{5}$$

where one of the two octants should be considered to generate data and the other octant should be used to simulate the theoretical model. From the expression of $P^{4\nu}_{\mu e}$, ΔP can be written as, $\Delta P = \Delta P_0 + \Delta P_1 + \Delta P_2$.

ΔP_0 is positive-definite. ΔP_1 and ΔP_2 depends on the CP-phases and can be both positive or negative. These terms can be expressed as

$$\Delta P_0 \simeq 8 \eta s^2_{13} \sin^2 \Delta \,. \tag{6}$$

$$\Delta P_1 = 4 s_{13} s_{12} c_{12} (\alpha \Delta) \sin \Delta \big[\cos(\Delta \pm \phi^{\rm HO}) - \cos(\Delta \pm \phi^{\rm LO}) \big] \,, \tag{7}$$

$$\Delta P_2 = 2\sqrt{2} s_{14} s_{24} s_{13} \sin \Delta \big[\sin(\Delta \pm \psi^{\rm HO}) - \sin(\Delta \pm \psi^{\rm LO}) \big] \,, \tag{8}$$

In above, $\phi = \delta_{13}$ and $\psi = \delta_{13} - \delta_{14}$ with the appropriate superscripts LO or HO. The term η is a positive definite angle and dictates the deviation from maximal mixing as $\theta_{23} = \frac{\pi}{4} \pm \eta$, where +(-) corresponds to HO (LO). If we need to measure the octant of θ_{23}, the contribution coming from ΔP_0 must not get cancelled completely in cases where the sum of ΔP_1 and ΔP_2 gives a negative contribution.

Results and discussion: Simulations for DUNE have been performed considering a total 248 kt.MW.yr of exposure, divided equally between ν and $\bar{\nu}$ mode. In fig.1, we show the bi-event plot. The ellipses (colored blobs) correspond to the 3ν (3+1 scheme) where, $\sin^2 \theta_{23} = 0.42\,(0.58)$ has been assumed as a benchmark value for the LO (HO). Since mass hierarchy can be measured relatively easily[‖] in DUNE, we can only concentrate on one of the two hierarchies,

[‖] In fig.1, we notice an small overlap between normal hierarchy (NH) and inverted hierarchy (IH) blobs which can be eliminated using the spectral information available in DUNE (see [8]).



Fig. 1: Bi-event plot for DUNE. The ellipses (colored blobs) corresond to the 3ν (3+1 scheme). $\sin^2\theta_{23} = 0.42\,(0.58)$ is taken as benchmark value for LO (HO). In cases of ellipses (colored blobs), the running parameter(s) is δ_{13} (δ_{13} & δ_{14}). In the 3+1 case, $\theta_{34} = 0^0$ has been assumed. This figure has been taken from [10].

say normal hierachy. While going from 3ν to 3+1 scheme, the ellipses becomes blobs because of the convolution of different combinations of δ_{13} & δ_{14} (see [8,9]). In this figure, we see a substantial overlap between LO and HO blobs due to the presence of the term ΔP_2, which depends on the new CP-phase δ_{14}. For detailed discussion, see [10].

Fig.2 depicts the discovery reach of θ_{23} octant in $\left[\sin^2\theta_{23}, \delta_{13}\right]$(true) plane assuming NH as true choice. Left (right) panel shows the results for 3ν (3+1) scheme. In 3ν case, a minimum 2σ sensitivity can be achieved if $\sin^2\theta_{23}(\text{true}) \lesssim 0.47$ and $\sin^2\theta_{23}(\text{true}) \gtrsim 0.55$ irrespective of the choice of δ_{13}(true). But, in 3+1 case we hardly have any octant sensitivity in the entire $\left[\sin^2\theta_{23}, \delta_{13}\right]$(true) plane.

Conclusions: In this work, we have studied the impact of a light eV-scale sterile neutrino in measuring the octant of θ_{23} at DUNE. The sensitivity towards θ_{23} octant can be completely lost if there is active- sterile oscillations.

Acknowledgments: S.S.C. would like to thank the organizers of XXII DAE-BRNS HEP Symposium 2016 for giving an opportunity to present this work. S.K.A. is supported by the DST/INSPIRE Research Grant [IFA-PH-12], Department of Science & Technology, India. A.P. is supported by the grant "Future In Research" *Beyond three neutrino families*, Fondo di Sviluppo e Coesione 2007-2013, APQ Ricerca Regione Puglia, Italy, Programma regionale a sostegno della specializzazione intelligente e della sostenibilità sociale ed ambientale. A.P. acknowledges partial support by the research project *TAsP* funded by the Instituto Nazionale di Fisica Nucleare (INFN).



Fig. 2: Discovery potential for excluding the wrong octant in $[\sin^2\theta_{23}, \delta_{13}]$ (true) plane assuming NH as true choice. The left (right) panel corresponds to the 3ν (3+1) case. In 3-flavor scenario, we marginalize over $(\theta_{23}, \delta_{13})$ (test). In 3+1 case, in addition, we marginalize over δ_{14} (true) and δ_{14} (test) fixing $\theta_{14} = \theta_{24} = 9^0$ and $\theta_{34} = 0$. This figure has been taken from [10].

References

1. G.L. Fogli and E. Lisi: Tests of three flavor mixing in long baseline neutrino oscillation experiments. Phys. Rev. D 54, 3667 (1996) [hep-ph/9604415]
2. Sanjib Kumar Agarwalla, Suprabh Prakash, and S.Uma Sankar: Resolving the octant of θ_{23} with T2K and NOνA. JHEP 1307 (2013) 131 [arXiv: 1301.2574]
3. K.N. Abazajian (UC, Irvine) et al.. : Light Sterile Neutrinos: A White Paper. [arXiv: 1204.5379]
4. N. Klop and A. Palazzo: Imprints of CP violation induced by sterile neutrinos in T2K data. Phys. Rev. D91 (2015) no.7, 073017 [arXiv: 1412.7524]
5. F. Capozzi, E. Lisi, A. Marrone, D. Montanino, and A. Palazzo: Neutrino masses and mixings: Status of known and unknown 3ν parameters. Nucl. Phys. B908, 218 (2016) [arXiv: 1601.07777]
6. C. Giunti, M. Laveder, Y.F.Li, and H.W.Long: Pragmatic view of Short-Baseline Neutrino Oscillations. Phys. Rev. D88, 073008 (2013)
7. J. Kopp, P.A.N. Machado, M. Maltoni, and T. Schwetz: Sterile Neutrino Oscillatios: The Global Picture. JHEP 1305 (2013) 050
8. Sanjib Kumar Agarwalla, Sabya Sachi Chatterjee and Antonio Palazzo: Physics Reach of DUNE with a Light Sterile Neutrino. JHEP 1609 (2016) 016 [arXiv: 1603.03759]
9. Sanjib Kumar Agarwalla, Sabya Sachi Chatterjee, Arnab Dasgupta and Antonio Palazzo: Discovery Potential of T2K and NOvA in the Presence of a Light Sterile Neutrino. JHEP 1602 (2016) 111 [arXiv: 1601.05995]
10. Sanjib Kumar Agarwalla, Sabya Sachi Chatterjee and Antonio Palazzo: Octant of θ_{23} in danger with a light sterile neutrino. Phys.Rev.Lett. 118 (2017) no.3, 031804 [arXiv: 1605.04299]